

# **2025 Volume 6** Issue 1 Edoc GF30089C7



## Spike Packet Coding: Lessons from Electric Fish\*

Ángel Ariel Caputi†

#### **Abstract**

Spike packet code is one of the least explored in the brain. Packet coding is based on three principles: 1) Sensory flow is composed of a series of discrete self-generated sensory image whose precise timing is separately encoded in the brain. 2) Images are encoded in the temporal structure of spike evoked trains, defined by onset time, inter-spike intervals, and spike number. 3) Packet information should be stored in some manner that allows memories to operate with subsequent sensory inputs. This form of coding may facilitate neural computations underlying natural behaviors, encompassing aspects such as novelty detection and boundary recognition. This article reviews some contributions from weakly electric fish that have advanced both the experimental and theoretical understanding of the spike packet neural code.

## **Keyphrases**

Spike timing, synaptic plasticity, active sensing.

#### Introduction

Ocular micro-saccades are followed by short-latency cortical evoked potentials (Gaarder et al. 1964). Gaarder (1966) posited that these potentials are used for the detection of borders, coining the phrase "packet information transmission". Under this hypothesis, the reafferent consequences of self-generated shifts of retinal images are incorporated in visual signals, thereby fragmenting the continuous stream of light into series of discrete and compact units (packets) of information that can be encoded, transmitted, stored, and manipulated. This notion was further substantiated by subsequent research involving monkeys freely looking at natural visual stimuli. Fixation-related spike synchronization occurs at the early phase of a rate response in some neurons of the primary visual cortex (Maldonado et al. 2008; Ito et al. 2011). This suggests that the oculomotor command might drive a corollary signal enabling precise timing of the earliest spike after every saccade. More recently, the concept of packet information transmission has been generalized and applied to the cortical processing of various sensory modalities, including audition, olfaction, and somatosensation (Luczak et al. 2015). The conceptualization of information packaging as spike trains carries significant theoretical implications: 1) The stream of information is segmented by precisely timed, discrete, and transient self-generated actions. 2) Individual images are encoded in packets consisting of spike timing probability distributions after the self-generated action. 3) Packet information may be retained in some manner that allows memories to operate with subsequent sensory inputs. In this article, I propose that these memories may be supported by temporally precise alterations in synaptic weights which work together as packet operands that process the next neuronal input to the same neuron. I present empirical evidence that substantiates this hypothesis within cerebellum-like sensory networks (the electrosensory lobes, ELs), typically present in two analogous yet non-homologous active electrosensory systems, and I elaborate on how the interplay of self- generation, spike packet encoding, storage, and operational mechanisms orchestrates novelty detection.

## Pulsatile electric discharges carry discrete selfgenerated electric images

African Mormyriformes and American Gymnotiformes emit electric organ discharges (EODs) converging into two main evolutionary basins: some species generate sinewave-like continuous electric fields while other generate series of brief pulses showing a species-specific time course separated by silences (wave and pulse fishes, respectively, Fig. 1). In the case of pulse fishes each EOD provokes the polarization of nearby




Figure 1: Species specific electric organ discharges (EODs) Four species belonging either to African and American taxa and exhibiting either wave or pulse electric discharges (adapted from Á. A. Caputi (2017)).

objects which, in turn, behave as virtual sources projecting electrosensory images on a cutaneous mosaic of electroreceptors. Here I focus on two paradigmatic species of pulse fish *Gymnotus omarorum* and

<sup>\*</sup>Report presented 2025-06-04, Neural Coding 2025, 16th International Neural Coding Workshop, Ascona Switzerland.

<sup>&</sup>lt;sup>†</sup>Correspondence to caputiangel@gmail.com.

Gnathonemus petersii.

## Sensory images are encoded as packets of post-EOD spike firing probabilities

The electrosensory mosaics of African pulse fish (Bell 1989; von der Emde and Bleckmann 1992) and American pulse fish (Á. A. Caputi and Aguilera 2019) comprise clusters of receptors specifically tuned to the typical time course of the species' EOD. These electroreceptors are innervated by primary afferent neurons that transduce and encode the temporal profile and amplitude of the local transcutaneous field into a burst of spikes (Fig. 2, bottom). The alteration in the time course (Fig. 2, color insets) or in the root mean square (rms) value of an EOD in the series elicits a well-defined behavior characterized by an abrupt and pronounced reduction of a few inter-EOD intervals followed by a gradual reversion to the preceding baseline (behavioral novelty response, BNR). In both taxa, these primary afferents project onto a laminar cerebellum-




Figure 2: Peripheral encoding of the time course of the stimuli. Changes in the time course of the EOD maintaining stimulus intensity (insets) alters the spiking pattern of the primary afferent (bottom) and elicits a BNR (top, adapted from Borde and Á. A. Caputi (2025)).

like network localized in the lower brainstem. These electrosensory lobes (ELs) are extensively interconnected with the contralateral EL and with neighboring praeminentialis nuclei (PNs), collectively forming a complex for early processing of electrosensory signals (Bell and Maler 2005). The downstream target is the torus semicircularis (TS), which serves as a crucial sensory hub participating in the regulation of electro-motor, skeletal-motor, and intricate behavioral responses. In G. omarorum, peri-EOD histograms of EL neurons firing in the absence of external objects reveal that deeply located neurons exhibit a sharp firing pattern approximately 10 ms after the main peak of the EOD (Fig 3A and B), whereas superficially located neurons show a delayed and more dispersed probability distribution (Fig 3C and D). Deep and superficial neurons display two distinct types of responses to increases in stimuli, "center on" and "center off". These resemblances have prompted their functional classification in wave fish (Clarke et al. 2015). Typically, in all neuron types, peri-EOD histograms show a silence spanning between 6 to 9 ms after the EOD (Fig. 3 gray bar, Pereira et al. (2014); Rodríguez-Cattáneo et al. (2024). This silence can be attributed to a significant inhibition elicited by the primary afferent volley. This inhibition is partly mediated by large multipolar neurons that project onto the basilar dendrites of both deep and superficial "on neurons", accompanied by an augmentation in the activity of interneurons that in-




Figure 3: Post-EOD firing patterns in *G. omarorum*. Note a) the blanked interval corresponding to the EOD and b) the lack of firing in all histograms between 6 and 9 ms (gray bar).

hibit "off neurons" (Berman and Maler 1998). Inhibition of "on neurons" typically attenuates - or completely blocks - the excitatory synaptic effects of the primary afferent volley along the basilar dendritic trunks. Deep "center on" neurons, which lack an apical dendritic tree, originate a feed-forward functional pathway. They project onto a subset of PN neurons, which in turn project to the eminentia granularis posterior, where they activate the granule neurons that give rise to the parallel fibers driving the distal branches of the apical trees of superficial neurons. Image processing in the EL of *G. petersii* shows notable similarities. Strikingly, in addition to the hindbrain circuitry, African species show an EOD command corollary discharge (EOCD) which is extensively distributed throughout the brainstem, including both the ELs and the PNs. There are two EOCD components (Bell 1989): The gating EOCD acts on deeply located neurons and facilitates the response of "center on" neurons to the EOD (Fig 4A). The plastic EOCD consist of a synaptic pattern elicited by the activation of a data-bus of parallel fibers and elicit synaptic patterns in which excitatory and inhibitory post-synaptic potentials counterbalance according to the changes in their relative weights (Bell, A. Caputi, and Grant 1997).

## Spike packets can be stored as and operated with synaptic expectations

Figure 4B illustrates the increase in late excitatory EOCD synaptic potentials after pairing the EOCD with peripheral stimulus evoking a large inhibition in a "center off neuron" of *G. petersii* (Bell, A. Caputi, Grant, and Serrier 1993; Bell, A. Caputi, and Grant 1997). This anti-Hebbian plastic effect is also elicited when the EOCD is paired with intracellular stimuli administered at varying delays (Fig. 4C). The interpretation via Occam's razor suggests that prior neuronal spiking activity is preserved as a modulation of the EOCD synaptic potentials occurring at spike-specific timing. Strikingly, as the data bus conveys a vast temporal array of activities to all traversed apical dendritic arbors, the pattern of spike timing probabilities is deftly reflected in the synaptic potential profile. Essentially, an 'image in negative', a counterpart of the input packet is retained, prepared to engage with the forthcoming spiking pattern. This phenomenon is particularly significant in certain interneu-




Figure 4: Corollary discharges in *G. petersii*. In A and B the EOD was suppressed. A) "Center on" neuron responding to an electrosensory stimulus applied at long (top) and short (bottom) delays after the EOD. B) Response to EOCD of a "center off" neuron. Top: in the absence of stimulus. Middle: while an electrical stimulus was applied at the center of the neuron receptive field. Bottom: in the absence of stimuli after a period of stimulation (dotted line represents control response) C) Evolution of the EOCD synaptic potentials when pairing with intracellular stimuli applied at different delays in the same neuron as in B (adapted from Bell, A. Caputi, and Grant (1997)).




Figure 5: Anti-Hebbian spike timing dependent plasticity. The top and third row respectively represent the EOCD synaptic potentials before and after intracellularly evoking a dendritic spike (second row) either before (left column), during (middle column) and after (right column). The EOD was suppressed the remaining neural command (bottom row) was used to trigger the stimulus (adapted from Bell, A. Caputi, and Grant (1997)).

rons, where the temporal coincidence between synaptic activity and dendritic spiking induces synaptic depression, while timing discrepancies either before or after coincidence foster synaptic potentiation.

This "Mexican hat time pattern" of the plastic mechanism, augments temporal precision, thereby optimizing packet storage (Fig. 5). This particular form of plasticity, first shown in preparations in vivo of *G. petersii* (Bell, A. Caputi, Grant, and Serrier 1993; Bell, A. Caputi, and Grant 1997), is currently designated as "spike time dependent plasticity". In vitro studies indicate that the temporal architecture of the "Mexican hat" plastic adaptation is realized through the synergistic interplay of a synapse-specific mechanism and a non-specific mechanism (Grant et al. 1996; Bell, Han, et al. 1997). Subsequent research the cerebral cortex of mammals have revealed analogous plastic phenomena, albeit exhibiting a Hebbian induction rule (Markram et al. 1997).

### Role of spike packet code in novelty detection

Behavioral experiments conducted on *G. petersii*, wherein the EOD was suppressed and replaced by an artificial EOD administered at varying delays post-command while preserving all other EOD parameters suggested that anti-Hebbian plasticity is involved in novelty detection (Hall et al. 1995). The relationship between novelty detection at the ELs and BNRs was explored in more detail in *G. omarorum*. In this species, comparable BNRs of the same amplitude are provoked by equivalent augmentations in EOD amplitude, yet they are elicited by singular deviant stimulus or under a step-and-hold paradigms (Fig. 6 A and B). In both cases, the amplitude of the BNR (BRNa) is a logarithmic function of the increase in the initial deviant EOD (Á. A. Caputi, Aguilera, and Castelló 2003). When the baseline stimulus is longer than 30 s the




Figure 6: Behavioral novelty responses. A) Experimental schematics. B) Novelty responses elicited by step and hold and deviant stimulus. C and D) BNR amplitude increases with stimulus intensity and the scaling constant positively correlates with baseline duration and number of images (adapted from Á. A. Caputi, Aguilera, and Castelló (2003)).

amplitude of the BNR is independent of the duration stimulus baseline. However, when the stimulation pattern comprised duty cycles of a constant period consisting of two alternating terms — one characterized by low object impedance (baseline) and the other by high object impedance (test) — the amplitude of the response was contingent upon the duration of the baseline term (Fig. 6C, orange and cyan symbols). Furthermore, the rate of increase (Fig. 6C, fitting lines) is a function of the number of EODs encompassed within the baseline term (Fig. 6D).

Simulation of the electro-sensorimotor loop fits the experimental results. Novelty detection at the EL was conceptualized as a first-order concentrated parameter system. In biological terms, the output of the EL compares the present afferent input with an expectation constructed by leaky integrating preceding images (Á. A. Caputi, Rodríguez-Cattáneo, et al. 2023; Á. A. Caputi, Waddell, et al. 2023). Confirming




Figure 7: BNRa is predicted by the amplitude of the novelty potential A) Responses to different strength of the stimulus, BNR (brown) novelty potential (blue). A) Responses to different strength of the stimulus, B) Responses to different stimulation duty cycles, C) Correlation between BNR and novelty potential at the EL (adapted from Á. A. Caputi, Waddell, et al. (2023). B) Correlation between BNR and novelty potential at the EL C) Responses to different stimulation duty cycles (adapted from Á. A. Caputi, Waddell, et al. (2023)).

this hypothesis, local field potential recordings in the EL (LFP) show that the first stimulus deviant from a constant amplitude baseline evoked a local novelty potential which amplitude predicted the amplitude of the BNR either when the stimulus amplitude or the baseline duration was manipulated (Figure 7, Á. A. Caputi, Waddell, et al. (2023)).

The present hypothesis posits that the time delay instantiated by the feed-forward pathway in *G. omarorum* conveys the expectation signal synaptic at the right moment to be compared with the afferent input. First, primary afferents activate excitatory contacts with thw basilar dendritic branches of "on neurons". Second, this excitatory input is counterbalanced by inhibitory projections from the ipsi- and contralateral deep inhibitory neurons, resulting in the typical silence that precedes the onset of the spike packet (Berman and Maler 1998). Third, "on neurons" receive a functional feed-forward through an EL-PN-EL pathway (Bastian and Courtright 1991).

This pathway originates from the projection of deep basilar neurons of the EL onto some PN neurons which subsequently project on a mass of granule cells dorsal to the EL. The axons of these granules form the typical cerebellum-like parallel fiber bus that activate the apical dendritic trees of superficial output neurons of the EL. Consequently, "on neurons" operate as comparators, integrating the afferent excitation of their basilar dendritic trees with the synaptic inputs on the apical dendritic trees. This signal integration is modulated by a feedback TS-PN pathway that project on the stems of the apical dendritic trees (Bastian, Chacron, et al. 2004).

Although definitive evidence for anti-Hebbian plasticity in the apical dendritic tree of "on neurons" in *G. omarorum* remains elusive, such phenomena have been documented in wave *Gymnotiformes* (Bastian 1999). Anti-Hebbian spike time-dependent synaptic plasticity within the apical dendritic arbor may modulate the intricate interplay between current inputs conveyed through afferent pathways and recent historical inputs preserved as modifications in synaptic efficacy, alongside the regulation of apical-somatic coupling. This adaptive filtering mechanism may enable the EL-PN circuit to attenuate stable components of

reafferent signals as well as other predictable patterns of sensory input, all while preserving responses to novel stimuli.

#### **Citation**

Brainiacs 2025 Volume 6 Issue 1 Edoc GF30089C7 Title: "Spike Packet Coding: Lessons from Electric Fish" Authors: Ángel Ariel Caputi

Dates: created 2025-05-01, presented 2025-06-04, updated 2025-

08-25, published 2025-08-31, revised 2025-09-10 Copyright: © 2025 Brain Health Alliance

Contact: caputiangel@gmail.com

NPDS: LINKS/Brainiacs/Caputi2025SPCLEF

DOI: 10.48085/GF30089C7

#### **Affiliations**

Ångel Ariel Caputi, caputiangel@gmail.com; Sistema Nacional de Investigadores, W. Ferreira Aldunate 1219 Pando, CP 15600 Montevideo, Uruguay.

#### References

- [1] J. Bastian. "Plasticity of feedback inputs in the apteronotid electrosensory system." *Journal of Experimental Biology* 202.10 (May 1999), pp. 1327–1337. ISSN: 1477-9145. DOI: 10.1242/jeb.202.10.1327 (cited p. 4).
- [2] J. Bastian, M. J. Chacron, and L. Maler. "Plastic and Nonplastic Pyramidal Cells Perform Unique Roles in a Network Capable of Adaptive Redundancy Reduction." *Neuron* 41.5 (Mar. 2004), pp. 767–779. ISSN: 0896-6273. DOI: 10.1016/s0896-6273(04)00071-6 (cited p. 4).
- [3] J. Bastian and J. Courtright. "Morphological correlates of pyramidal cell adaptation rate in the electrosensory lateral line lobe of weakly electric fish." *Journal of Comparative Physiology A* 168.4 (Apr. 1991), pp. 393–407. ISSN: 1432-1351. DOI: 10.1007/bf00199600 (cited p. 4).
- [4] C. C. Bell, A. Caputi, K. Grant, and J. Serrier. "Storage of a sensory pattern by anti-Hebbian synaptic plasticity in an electric fish." *Proceedings of the National Academy of Sciences* 90.10 (May 1993), pp. 4650–4654. ISSN: 1091-6490. DOI: 10.1073/pnas.90.10.4650 (cited pp. 2, 3).
- [5] C. C. Bell. "Sensory coding and corollary discharge effects in mormyrid electric fish." *Journal of Experimental Biology* 146.1 (Sept. 1989), pp. 229– 253. ISSN: 1477-9145. DOI: 10.1242/jeb.146.1.229 (cited p. 2).
- [6] C. C. Bell and L. Maler. "Central neuroanatomy of electrosensory systems in fish." In: *Electroreception*. New York: Springer, 2005, pp. 68–111 (cited p. 2).
- [7] C. C. Bell, A. Caputi, and K. Grant. "Physiology and Plasticity of Morphologically Identified Cells in the Mormyrid Electrosensory Lobe." *The Journal of Neuroscience* 17.16 (Aug. 1997), pp. 6409–6423. ISSN: 1529-2401. DOI: 10.1523/jneurosci.17-16-06409.1997 (cited pp. 2, 3).
- [8] C. C. Bell, V. Z. Han, Y. Sugawara, and K. Grant. "Synaptic plasticity in a cerebellum-like structure depends on temporal order." *Nature* 387.6630 (May 1997), pp. 278–281. ISSN: 1476-4687. DOI: 10.103 8/387278a0 (cited p. 3).

- [9] N. J. Berman and L. Maler. "Inhibition Evoked From Primary Afferents in the Electrosensory Lateral Line Lobe of the Weakly Electric Fish (Apteronotus leptorhynchus)." *Journal of Neurophysiology* 80.6 (Dec. 1998), pp. 3173–3196. ISSN: 1522-1598. DOI: 10.1152/jn.1998.80 .6.3173 (cited pp. 2, 4).
- [10] M. Borde and Á. A. Caputi. "The sensory-effector cycle, contributions from a native species." Neuroscience 571 (Apr. 2025), pp. 108–129. ISSN: 0306-4522. DOI: 10.1016/j.neuroscience.2025.02.015 (cited p. 2).
- Á. A. Caputi. "The bioinspiring potential of weakly electric fish." Bioinspiration & Biomimetics 12.2 (Feb. 2017), p. 025004. ISSN: 1748-3190.
  DOI: 10.1088/1748-3190/12/2/025004 (cited p. 1).
- [12] Á. A. Caputi, P. A. Aguilera, and M. E. Castelló. "Probability and amplitude of novelty responses as a function of the change in contrast of the reafferent image inG. carapo." *Journal of Experimental Biology* 206.6 (Mar. 2003), pp. 999–1010. ISSN: 0022-0949. DOI: 10.1242/jeb.00199 (cited p. 3).
- [13] Á. A. Caputi and P. A. Aguilera. "Encoding phase spectrum for evaluating "electric qualia"." *Journal of Experimental Biology* (Jan. 2019). ISSN: 0022-0949. DOI: 10.1242/jeb.191544 (cited p. 2).
- [14] Á. A. Caputi, A. Rodríguez-Cattáneo, J. C. Waddell, A. C. Pereira, and P. A. Aguilera. "Getting the news in milliseconds: The role of early novelty detection in active electrosensory exploration." *Biosystems* 223 (Jan. 2023), p. 104803. ISSN: 0303-2647. DOI: 10.1016/j.biosystems.2022.104803 (cited p. 4).
- [15] Á. A. Caputi, J. C. Waddell, and P. A. Aguilera. "A simple model of the electrosensory electromotor loop in Gymnotus omarorum." *Biosystems* 223 (Jan. 2023), p. 104800. ISSN: 0303-2647. DOI: 10.1016/j.biosystems.2022.104800 (cited p. 4).
- [16] S. E. Clarke, A. Longtin, and L. Maler. "Contrast coding in the electrosensory system: parallels with visual computation." *Nature Reviews Neuro*science 16.12 (Nov. 2015), pp. 733–744. ISSN: 1471-0048. DOI: 10.103 8/nrn4037 (cited p. 2).
- [17] K. Gaarder. "Transmission of Edge Information in the Human Visual System." Nature 212.5059 (Oct. 1966), pp. 321–323. ISSN: 1476-4687. DOI: 10.1038/212321a0 (cited p. 1).
- [18] K. Gaarder, J. Krauskopf, V. Graf, W. Kropfl, and J. C. Armington. "Averaged Brain Activity Following Saccadic Eye Movement." Science 146.3650 (Dec. 1964), pp. 1481–1483. ISSN: 1095-9203. DOI: 10.1126/science.146.3650.1481 (cited p. 1).
- [19] K. Grant, C. Bell, and V. Han. "Sensory expectations and anti-Hebbian synaptic plasticity in cerebellum-like structures." *Journal of Physiology-Paris* 90.3–4 (Jan. 1996), pp. 233–237. ISSN: 0928-4257. DOI: 10.101 6/s0928-4257 (97) 81430-1 (cited p. 3).
- [20] C. Hall, C. Bell, and R. Zelick. "Behavioral evidence of a latency code for stimulus intensity in mormyrid electric fish." *Journal of Comparative Physiology A* 177.1 (July 1995). ISSN: 1432-1351. DOI: 10.1007/bf002 43396 (cited p. 3).
- [21] J. Ito, P. Maldonado, and S. Gruen. "Cross-frequency coupling of eye-movement related LFP activities of freely viewing monkeys." BMC Neuroscience 12.S1 (July 2011). ISSN: 1471-2202. DOI: 10.1186/1471-2202-12-s1-p132 (cited p. 1).
- [22] A. Luczak, B. L. McNaughton, and K. D. Harris. "Packet-based communication in the cortex." *Nature Reviews Neuroscience* 16.12 (Oct. 2015), pp. 745–755. ISSN: 1471-0048. DOI: 10.1038/nrn4026 (cited p. 1).

- P. Maldonado, C. Babul, W. Singer, E. Rodriguez, D. Berger, and S. Grün. "Synchronization of Neuronal Responses in Primary Visual Cortex of Monkeys Viewing Natural Images." *Journal of Neurophysiology* 100.3 (Sept. 2008), pp. 1523–1532. ISSN: 1522-1598. DOI: 10.1152/jn.00076.2008 (cited p. 1).
- [24] H. Markram, J. Lübke, M. Frotscher, and B. Sakmann. "Regulation of Synaptic Efficacy by Coincidence of Postsynaptic APs and EPSPs." Science 275.5297 (Jan. 1997), pp. 213–215. ISSN: 1095-9203. DOI: 10.112 6/science.275.5297.213 (cited p. 3).
- [25] A. C. Pereira, A. Rodríguez-Cattáneo, and A. A. Caputi. "The slow pathway in the electrosensory lobe of Gymnotus omarorum: Field potentials and unitary activity." *Journal of Physiology-Paris* 108.2–3 (Apr. 2014), pp. 71–83. ISSN: 0928-4257. DOI: 10.1016/j.jphysparis.2014.07.005 (cited p. 2).
- [26] A. Rodríguez-Cattáneo, A. C. Pereira, P. A. Aguilera, and Á. A. Caputi. "Packet information encoding in a cerebellum-like circuit." *PLOS ONE* 19.9 (Sept. 2024). Ed. by G. S. Cymbalyuk, e0308146. ISSN: 1932-6203. DOI: 10.1371/journal.pone.0308146 (cited p. 2).
- [27] G. von der Emde and H. Bleckmann. "Differential responses of two types of electroreceptive afferents to signal distortions may permit capacitance measurement in a weakly electric fish, Gnathonemus petersii." *Journal of Comparative Physiology A* 171.5 (Sept. 1992), pp. 683–694. ISSN: 1432-1351. DOI: 10.1007/bf00194116 (cited p. 2).